

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



WAX BLOCKS



RODENTICIDE

Active Constituent: 0.05 g/kg BRODIFACOUM

Ready to use block bait for use in damp or dry situations in and around industrial, commercial, public services, agricultural and domestic buildings.

One feed kills rats and mice.

NRA Approval No: 52675/0202

Pack size: 1 kg

GENERAL INSTRUCTIONS

Locate bait stations along walls or rafters and in dark sheltered spots where there are signs of rodent activity. Nail in place if relevant. Wax blocks are particularly suitable for use on damp situations and sewers, and crop stores where the risk of contamination should be minimised. Record the number and location of bait stations for the purpose of replenishment and later removal. Most deaths occur 4 to 7 days after consumption of bait.

Note: One feed is sufficient to kill rats and mice. Rats and mice will continue to eat the bait after a lethal dose has been eaten.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Depending on the amount eaten, this bait may kill or injure. The bait is hazardous to dogs, cats, pigs, poultry and other wildlife. DO NOT place baits in locations which are accessible to pets, domestic animals, livestock or birds. DO NOT contaminate dams, waterways and drains with the bait or its used container. The use of this product for the control of protected native rodents and rodent like animals requires permission from wildlife authorities. The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces, e.g. drains.

STORAGE AND DISPOSAL

Containers which have held bait should not be used for any other purpose. Store in the closed original container in a dry place out of the reach of children. DO NOT store in direct sunlight. Dispose of empty containers and unused or untaken baits by burying in an approved dump. Look for dead rats or mice, burn or bury them, DO NOT place in refuse bins or rubbish tips. Ensure baits are removed when rodent activity has ceased.

SAFETY DIRECTIONS

Poisonous if swallowed. DO NOT touch bait. If on skin and after each baiting, wash thoroughly with soap and water. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126. Vitamin K1 (Phytomenadione) is antidotal.

NOTE TO PHYSICIANS AND VETERINARIANS

Brodifacoum, the active constituent of TALON Wax Blocks, is a long lasting anticoagulant chemical, which if ingested by humans, domestic animals or pets, can reduce the clotting power of the blood and haemorrhage may result. Vitamin K1 (Phytomenadione) should be administered urgently by medical, appropriate para-medical personnel or veterinarians. If administered intravenously the injection must be given slowly.

MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet. For a copy phone 1800 067 108 or visit our website at www.syngenta.com.au

MANUFACTURER'S WARRANTY AND EXCLUSION OF LIABILITY

Syngenta Crop Protection Pty Limited has no control over storage, handling and manner of use of this product. Where this material is not stored, handled or used correctly and in accordance with directions, no express or implied representations or warranties concerning this product (other than non-excludable statutory warranties) will apply. Syngenta accepts no liability for any loss or damage arising from incorrect storage, handling or use.

® Registered trademark of a Syngenta Group Company.

* Bitrex is a trademark of Macfarlan Smith Limited.

DIRECTIONS FOR USE

Restrictions: DO NOT apply baits to crops.
 DO NOT place bait stations in the open.
 DO NOT exceed 3 m between bait stations for mice or 9 m for rats.

Baiting Strategy

- Eliminate as far as practical, all alternative food sources.
- Bait infested area. Place baits under cover. Inspect baits daily.

For rats: Use 1 to 3 blocks depending on the level of infestation.

For mice: Use single blocks.

- Replace eaten baits. If eaten quickly increase number of baits. Continue observation and replenishment until no more baits are taken. Baiting for at least 2 weeks will be necessary to reduce rat/mouse numbers.
- Ensure baits are removed when rodent activity has ceased.

Application (All States)

Situation	Pest	Rate	Critical Comments
Industrial, Commercial, Public Services, Agricultural and Domestic Buildings	Rats and mice (<i>Rattus rattus</i> , <i>R. norvegicus</i> , <i>Mus domesticus</i>) only.	1 to 3 blocks / bait station	Set out baits in places frequented by mice and rats but not more than 2 m away from buildings etc. For rats: Place one to three blocks at intervals of 5 to 9 m in infested areas. For mice: Place single blocks at 2 to 3 m intervals in infested areas.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.